



HORTI-TRACE MIX

FULLY CHELATED

MAGNESIUM 2% w/v | ZINC 2.5% w/v | MANGANESE 3% w/v
SULFUR 6.2% w/v | COPPER .5% w/v | IRON 1% w/v
BORON .04% w/v | MOLYBDENUM .05% w/v



CHELATED FOR SUPERIOR ABSORPTION AND QUICK UPTAKE OF NUTRIENTS

TYPICAL ANALYSIS	AMOUNT	UNITS
PH	3.8	units
Specific gravity	1.32	
Filtered	100	microns
Trace Elements		
Magnesium (Mg)	2.0	%w/v
Zinc (Zn)	2.5	%w/v
Manganese (Mn)	3.0	%w/v
Sulphur (S)	6.2	%w/v
Copper (Cu)	0.5	%w/v
Iron (Fe)	1.0	%w/v
Boron (B)	0.04	%w/v
Molybdenum (Mo)	0.05	%w/v

- ✓ Highly effective lignosulphonate chelating agent
- ✓ Superior absorption with rainfast properties

Kemgro™ Horti-trace Mix provides you with a complete balance of trace elements. Suitable for most horticultural crops. Chelated with a highly effective chelating agent for superior absorption, quick up take of nutrients with rainfast properties. Can be applied as a foliar spray or through a fertigation system. Filtered to 100 microns for easy application.

Note: Kemgro recommends Bowa Spray Activator Surfactant to be used with this product.

DIRECTIONS FOR USE Agitate well or stir before use. Always dilute and do not exceed recommended rate per hectare.

Horticulture Use at least 300L/ha of water.

CROP	RATE/HA	APPLICATION RATES AND TIMING
Vines	2-4 L/ha	At 10cm shoot, post-flowering and post-harvest.
Citrus	2-4 L/ha	At Spring and Autumn flush, and as required.
Stone & pome fruit	2-4 L/ha	Post-flowering and after harvest.
Avocado	2-3 L/ha	At Spring flush and post-harvest.
Vegetables	1-4 L/ha	When needed or with normal nutrient program.
Almonds	2-3 L/ha	At petal fall, 4-6 weeks later and post-harvest.
Potatoes	2-4 L/ha	At hooking and tuber set.
Soil drench	5-10 L/ha	Prior to planting.
Seed dressing	7-9 L	Per tonne of seed.
Fertigation	5-9 L/ha	As required, apply in the middle of irrigation to allow flushing.

AVAILABLE IN

