

and operated

KEMGRO ZMC ORGANIC FULLY CHELATED



CHELATED FOR SUPERIOR ABSORPTION AND QUICK UPTAKE OF NUTRIENTS

TYPICAL ANALYSIS	AMOUNT	UNITS
PH	3.8	units
Specific gravity	1.32	
Filtered	100	microns
Zinc (Zn)	6.3	%w/v
Manganese (Mn)	5.1	%w/v
Copper (Cu)	1.5	%w/v
Molybdenum (Mo)	0.1	%w/v
Sulphur (S)	6.9	%w/v
Sulphur (S)	6.9	%w/v



✓ Highly effective lignosulphonate chelating agent ✓ Superior absorption with rainfast properties ✓ Australian manufacturer

Kemgro[®] ZMC Organic nutrient is chelated with a highly effective chelating agent for superior absorption, quick uptake of nutrients with rainfast properties. Can be applied as a foliar spray or through a fertigation system. Filtered to 100 microns for easy application.

DIRECTIONS FOR USE Agitate well before use. Always use diluted and do not exceed recommended rate per hectare.

HORTICULTURE: Use at least 300L/ha of water. **BROADACRE**: Use at least 80L/ha of water.

CROP	RATE/HA	APPLICATION RATES AND TIMING	
Vines	2-4 L/ha	At 10cm shoot, post-flowering and post-harvest.	
Citrus	2-4 L/ha	At Spring and Autumn flush, and as required.	
Stone & pome fruit	2-4 L/ha	Post-flowering and after harvest.	
Avocado	2-3 L/ha	At Spring flush and post-harvest.	
Vegetables	1-4 L/ha	When needed or with normal nutrient program.	
Almonds	2-3 L/ha	At petal fall, 4-6 weeks later and post-harvest.	
Potatoes	2-4 L/ha	At hooking and tuber set.	
Seed dressing	7-9 L	Per tonne of seed.	
Fertigation	5-9 L/ha	As required, apply in the middle of irrigation to allow flushing.	
Seeding in furrow	2-4L/ha	Apply at planting.	
Broadacre	2-4L/ha	Apply when sufficient foliage is present. Spray as required.	

