



BORON

FULLY CHELATED

BORON 15% w/v | NITROGEN 6.9% w/v



- ✓ Highly effective lignosulphonate chelating agent
- ✓ Superior absorption with rainfast properties
- ✓ Local Australian Manufacturer

Kemgro[®] Boron nutrient is chelated with a highly effective chelating agent for superior absorption, quick uptake of nutrients with rainfast properties. Can be applied as a foliar spray or through a fertigation system. Filtered to 100 microns for easy application.

Note: Kemgro recommends Bowa Spray Activator Surfactant to be used with this product.

DIRECTIONS FOR USE Agitate well or stir before use. Always dilute and do not exceed recommended rate per hectare.

HORTICULTURE Use at least 300L/ha of water.

BROADACRE Use at least 80L/ha of water.

WARNING

Avoid spraying at flowering.

**CHELATED FOR SUPERIOR
ABSORPTION AND QUICK
UPTAKE OF NUTRIENTS**

TYPICAL ANALYSIS	AMOUNT	UNITS
PH	8.0	units
Specific gravity	1.38	
Filtered	100	microns
Major Elements		
Boron (B)	15.0	%w/v
Nitrogen (N)	6.9	%w/v

CROP	RATE/HA	APPLICATION RATES AND TIMING
Vines	1-3 L/ha	When clusters are visible, 2nd at flower bud separation.
Citrus	1-3 L/ha	At spring flush. A 2nd application may be required.
Stone & pome fruit	1-3 L/ha	Early fruit set. A 2nd application may be required.
Avocado	1-3 L/ha	Early fruit set. A 2nd application may be required.
Vegetables	1-3 L/ha	Before flowering. A 2nd application may be required.
Almonds	1-3 L/ha	Apply with 1st cover spray. A 2nd application may be required.
Potatoes	1-3 L/ha	1-2 applications during tuber growth.
Broadacre	1-2 L/ha	Mid to late tillering.
Soil drench	2-4 L/ha	Prior to planting.
Fertigation	2-5 L/ha	As required, apply in the middle of irrigation to allow flushing.

Frequencies are guidelines only.

AVAILABLE IN

