



CALCIUM

FULLY CHELATED

CALCIUM 15% w/v | NITROGEN 12% w/v



- ✓ Highly effective lignosulphonate chelating agent
- ✓ Superior absorption with rainfast properties
- ✓ Local Australian Manufacturer

Kemgro® Calcium nutrient is chelated with a highly effective chelating agent for superior absorption, quick uptake of nutrients with rainfast properties. Can be applied as a foliar spray or through a fertigation system. Filtered to 100 microns for easy application.

Note: Kemgro recommends Bowa Spray Activator Surfactant to be used with this product.

DIRECTIONS FOR USE Agitate well or stir before use. Always dilute and do not exceed recommended rate per hectare.

HORTICULTURE Use at least 300L/ha of water.

BROADACRE Use at least 80L/ha of water.

CHELATED FOR SUPERIOR ABSORPTION AND QUICK UPTAKE OF NUTRIENTS

| TYPICAL ANALYSIS | AMOUNT | UNITS |
|-----------------------|--------|---------|
| PH | 5.0 | units |
| Specific gravity | 1.43 | |
| Filtered | 100 | microns |
| Major Elements | | |
| Calcium (Ca) | 15.0 | %w/v |
| Nitrogen (N) | 12.0 | %w/v |

| CROP | RATE/HA | APPLICATION RATES AND TIMING |
|--------------------|------------|---|
| Vines | 3-5 L/ha | At bud burst, fruit set and 2 weeks later. |
| Citrus | 4-6 L/ha | Pre-flowering, fruit set and 4-6 weeks later. |
| Stone & pome fruit | 3-5 L/ha | At bud burst, fruit set and 2 weeks later. |
| Avocado | 3-5 L/ha | At bud burst, fruit set and 2 weeks later. |
| Vegetables | 2-4 L/ha | From 4-6 leaf 2-5 applications. |
| Potatoes | 4-6 L/ha | At hooking and tuber set. |
| Broadacre | 2-4 L/ha | When sufficient foliage is present spray as required. |
| Soil drench | 10-25 L/ha | Prior to planting. |
| Fertigation | 10-25 L/ha | As required, apply in the middle of irrigation to allow flushing. |

Frequencies are guidelines only.

AVAILABLE IN

