

KEMGRO®

COPPER

FULLY CHELATED

COPPER 6% w/v | SULFUR 3.1% w/v



- ✓ Highly effective lignosulphonate chelating agent
- ✓ Superior absorption with rainfast properties
- ✓ Local Australian Manufacturer

Kemgro® Copper nutrient is chelated with a highly effective chelating agent for superior absorption, quick uptake of nutrients with rainfast properties. Can be applied as a foliar spray or through a fertigation system. Filtered to 100 microns for easy application.

Note: Kemgro recommends Bowa Spray Activator Surfactant to be used with this product.

DIRECTIONS FOR USE Agitate well or stir before use. Always dilute and do not exceed recommended rate per hectare.

HORTICULTURE Use at least 300L/ha of water.

BROADACRE Use at least 80L/ha of water.

CHELATED FOR SUPERIOR ABSORPTION AND QUICK UPTAKE OF NUTRIENTS

TYPICAL ANALYSIS	AMOUNT	UNITS
PH	3.1	units
Specific gravity	1.15	
Filtered	100	microns
Major Elements		
Copper (Cu)	6.0	%w/v
Sulfur (S)	3.1	%w/v

CROP	RATE/HA	APPLICATION RATES AND TIMING
Vines	1-2 L/ha	Apply if a deficiency is evident.
Citrus	1-2 L/ha	At Spring flush. A second application may be required. Avoid spraying at blossom and harvest
Stone & pome fruit	1-2 L/ha	Apply if a deficiency is evident.
Avocado	1-2 L/ha	At Spring flush. A second application may be required.
Vegetables	1-2 L/ha	Before flowering. A second application may be required.
Almonds	1-2 L/ha	Apply with first cover spray. A second application may be required.
Potatoes	1-2 L/ha	Tuber set and prior to closure.
Broadacre	.5-1 L/ha	When sufficient foliage is present spray as required.
Soil drench	2-4 L/ha	Prior to planting.
Fertigation	3-6 L/ha	As required, apply in the middle of irrigation to allow flushing.

Frequencies are guidelines only.

AVAILABLE IN

