

# KEMGRO®

# IRON

## FULLY CHELATED

IRON 6% w/v | SULFUR 3.4% w/v



- ✓ Highly effective lignosulphonate chelating agent
- ✓ Superior absorption with rainfast properties
- ✓ Local Australian Manufacturer

Kemgro® Iron nutrient is chelated with a highly effective chelating agent for superior absorption, quick uptake of nutrients with rainfast properties. Can be applied as a foliar spray or through a fertigation system. Filtered to 100 microns for easy application.

Note: Kemgro recommends Bowa Spray Activator Surfactant to be used with this product.

**DIRECTIONS FOR USE** Agitate well or stir before use. Always dilute and do not exceed recommended rate per hectare.

**HORTICULTURE** Use at least 300L/ha of water.

**BROADACRE** Use at least 80L/ha of water.

**CHELATED FOR SUPERIOR ABSORPTION AND QUICK UPTAKE OF NUTRIENTS**

TYPICAL ANALYSIS	AMOUNT	UNITS
PH	2.1	units
Specific gravity	1.18	
Filtered	100	microns
<b>Major Elements</b>		
Iron (Fe)	6.0	%w/v
Sulfur (S)	3.4	%w/v

CROP	RATE/HA	APPLICATION RATES AND TIMING
Vines	1-3 L/ha	Apply if a deficiency is evident.
Citrus	1-3 L/ha	At Spring flush. A second application may be required. Avoid spraying at blossom and harvest
Stone & pome fruit	1-3 L/ha	Apply if a deficiency is evident.
Avocado	1-3 L/ha	At Spring flush. A second application may be required.
Vegetables	1-3 L/ha	Before flowering. A second application may be required.
Almonds	1-3 L/ha	Apply with first cover spray. A second application may be required.
Potatoes	1-3 L/ha	1-2 applications during tuber growth.
Broadacre	1-2 L/ha	When sufficient foliage is present spray as required.
Soil drench	2-6 L/ha	Prior to planting.
Fertigation	4-8 L/ha	As required, apply in the middle of irrigation to allow flushing.

*Frequencies are guidelines only.*

AVAILABLE IN

