KEMGRO MOLYBDENUM FULLY CHELATED

MOLYBDENUM 25% w/v | PHOSPHOROUS 11% w/v



CHELATED FOR SUPERIOR ABSORPTION AND QUICK UPTAKE OF NUTRIENTS

TYPICAL ANALYSIS	AMOUNTUNITS
PH	4.0 units
Specific gravity	1.58
Filtered	100 microns
Major Elements	
Molybdenum (Mo)	25.0 %w/v
Phosphorous (P)	11.0 %w/v



Highly effective lignosulphonate chelating agent Superior absorption with rainfast properties Local Australian Manufacturer

Kemgro[®] Molybdenum nutrient is chelated with a highly effective chelating agent for superior absorption, quick uptake of nutrients with rainfast properties. Can be applied as a foliar spray or through a fertigation system. Filtered to 100 microns for easy application.

Note: Kemgro recommends Bowa Spray Activator Surfactant to be used with this product.

DIRECTIONS FOR USE Agitate well or stir before use. Always dilute and do not exceed recommended rate per hectare.

HORTICULTURE Use at least 300L/ha of water. BROADACRE Use at least 80L/ha of water.

WARNING

Excessive use of Molybdenum can be harmful to stock. Users are encouraged to seek further advice.

CROP	RATE/HA	APPLICATION RATES AND TIMING
Vines	200-400 ml/ha	Apply early growth stage.
Citrus	200-400 ml/ha	At spring flush.
Stone & pome fruit	100-200 ml/ha	Apply when required.
Avocado	300-500 ml/ha	Apply when required.
Vegetables	150-300 ml/ha	At 4-5 leaf stage or as required.
Almonds	100-300 ml/ha	Apply when required.
Olives	100-300 ml/ha	Apply when required.
Potatoes	100-300 ml/ha	At early leaf stage or as required.
Broadacre	100-200 ml/ha	When sufficient foliage is present.
Fertigation	200-500 ml/ha	As required, apply in the middle of irrigation to allow flushing.
The second s		

Frequencies are guidelines only.

