



ZINC MANGANESE

FULLY CHELATED

ZINC 7.5% w/v | MANGANESE 7.5% w/v | SULFUR 8% w/v



- ✓ Highly effective lignosulphonate chelating agent
- ✓ Superior absorption with rainfast properties
- ✓ Local Australian Manufacturer

Kemgro[®] Zinc-Manganese nutrient is chelated with a highly effective chelating agent for superior absorption, quick uptake of nutrients with rainfast properties. Can be applied as a foliar spray or through a fertigation system. Filtered to 100 microns for easy application.

Note: Kemgro recommends Bowa Spray Activator Surfactant to be used with this product.

DIRECTIONS FOR USE Agitate well or stir before use. Always dilute and do not exceed recommended rate per hectare.

HORTICULTURE Use at least 300L/ha of water.

BROADACRE Use at least 80L/ha of water.

CHELATED FOR SUPERIOR ABSORPTION AND QUICK UPTAKE OF NUTRIENTS

TYPICAL ANALYSIS	AMOUNT	UNITS
PH	4.9	units
Specific gravity	1.4	
Filtered	100	microns
Major Elements		
Zinc (Zn)	7.5	%w/v
Manganese (Mn)	7.5	%w/v
Sulfur (S)	8.0	%w/v

CROP	RATE/HA	APPLICATION RATES AND TIMING
Vines	2-4 L/ha	At 10cm shoot, post-flowering and post-harvest.
Citrus	2-4 L/ha	At Spring and Autumn flush, and as required.
Stone & pome fruit	2-4 L/ha	Post-flowering and after harvest.
Avocado	2-3 L/ha	At Spring flush and post-harvest.
Vegetables	1-4 L/ha	When needed or with normal nutrient program.
Almonds	2-3 L/ha	At petal fall, 4-6 weeks later and post-harvest.
Potatoes	2-4 L/ha	At hooking and tuber set.
Broadacre	1-3 L/ha	When sufficient foliage is present spray as required.
Soil drench	5-10 L/ha	Prior to planting.
Seed dressing	7-9 L	Per tonne of seed.
Fertigation	5-9 L/ha	As required, apply in the middle of irrigation to allow flushing.

Frequencies are guidelines only.

AVAILABLE IN

