

## **FULLY CHELATED**

ZINC 4% w/v | MAGNESIUM 4% w/v | MANGANESE 4% w/v | NITROGEN 4.6% w/v | SULFUR 4.4% w/v



- ✓ Highly effective lignosulphonate chelating agent
- ✓ Superior absorption with rainfast properties
- ✓ Local Australian Manufacturer

Kemgro<sup>®</sup> Zinc-MG-MN nutrient is chelated with a highly effective chelating agent for superior absorption, quick uptake of nutrients with rainfast properties. Can be applied as a foliar spray or through a fertigation system. Filtered to 100 microns for easy application.

Note: Kemgro recommends Bowa Spray Activator Surfactant to be used with this product.

RATE/HA APPLICATION RATES AND TIMING

**DIRECTIONS FOR USE** Agitate well or stir before use. Always dilute and do not exceed recommended rate per hectare.

**HORTICULTURE** Use at least 300L/ha of water. **BROADACRE** Use at least 80L/ha of water.

## CHELATED FOR SUPERIOR ABSORPTION AND QUICK UPTAKE OF NUTRIENTS

TYPICAL ANALYSIS	AMOUNTUNITS	
PH	4.4	units
Specific gravity	1.37	
Filtered	100	microns
Major Elements		
Zinc (Zn)	4.0	%w/v
Magnesium (Mg)	4.0	%W/V
Manganese (Mn)	4.0	%w/v
Nitrogen (N)	4.6	%w/v
Sulfur (S)	4.4	%w/v

Vines	2-5 L/ha	At 10cm shoot, post-flowering and post-harvest.
Citrus	2-5 L/ha	At Spring and Autumn flush, and as required.
Stone & pome fruit	2-5 L/ha	Post-flowering and after harvest.
Avocado	2-4 L/ha	At Spring flush and post-harvest.
Vegetables	2-5 L/ha	When needed or with normal nutrient program.
Almonds	2-3 L/ha	At petal fall, 4-6 weeks later and post-harvest.
Potatoes	2-5 L/ha	At hooking and tuber set.
Broadacre	2-4 L/ha	When sufficient foliage is present spray as required.
Soil drench	5-10 L/ha	Prior to planting.
Seed dressing	7-9 L	Per tonne of seed.
Fertigation	5-15 L/ha	As required, apply in the middle of irrigation to allow flushing.
Frequencies are auidelines only.		

## **AVAILABLE IN**

