

# KEMGRO<sup>®</sup>

# ZINC

## FULLY CHELATED

ZINC 16% w/v | SULFUR 8.1% w/v



- ✓ Highly effective lignosulphonate chelating agent
- ✓ Superior absorption with rainfast properties
- ✓ Local Australian Manufacturer

Kemgro<sup>®</sup> Zinc nutrient is chelated with a highly effective chelating agent for superior absorption, quick uptake of nutrients with rainfast properties. Can be applied as a foliar spray or through a fertigation system. Filtered to 100 microns for easy application.

Note: Kemgro recommends Bowa Spray Activator Surfactant to be used with this product.

**DIRECTIONS FOR USE** Agitate well or stir before use. Always dilute and do not exceed recommended rate per hectare.

**HORTICULTURE** Use at least 300L/ha of water.

**BROADACRE** Use at least 80L/ha of water.

**CHELATED FOR SUPERIOR ABSORPTION AND QUICK UPTAKE OF NUTRIENTS**

TYPICAL ANALYSIS	AMOUNT	UNITS
PH	4.1	units
Specific gravity	1.4	
Filtered	100	microns
<b>Major Elements</b>		
Zinc (Zn)	16	%w/v
Sulfur (S)	8.1	%w/v

CROP	RATE/HA	APPLICATION RATES AND TIMING
Vines	2-4 L/ha	At 10cm shoot, post-flowering and post-harvest.
Citrus	2-4 L/ha	At Spring and Autumn flush, and as required.
Stone & pome fruit	2-4 L/ha	Post-flowering and after harvest.
Avocado	2-3 L/ha	At Spring flush and post-harvest.
Vegetables	1-4 L/ha	When needed or with normal nutrient program.
Almonds	2-3 L/ha	At petal fall, 4-6 weeks later and post-harvest.
Potatoes	2-4 L/ha	At hooking and tuber set.
Broadacre	1-3 L/ha	When sufficient foliage is present spray as required.
Soil drench	3-7 L/ha	Prior to planting.
Seed dressing	5-6 L	Per tonne of seed.
Fertigation	5-9 L/ha	As required, apply in the middle of irrigation to allow flushing.

*Frequencies are guidelines only.*

AVAILABLE IN

